

MAJOR GEOMAGNETIC STORMS AND COSMIC RAYS, 2. METHODS OF CR USING FOR FORECASTING.

A. V. Belov (1), L.I. Dorman (1,2), E. A. Eroshenko (1), R.T. Gushchina (1), N. Iucci (3), H. Mavromichalaki (4), L.A. PUSTIL'NIK (2), A. Sternlieb (2), G. Villoresi (3), V.G. Yanke (1), I.G. Zukerman (2)

(1) IZMIRAN, Russian Academy of Science, Troitsk; (2) Israel Cosmic Ray Center and Emilio Segre' Observatory, affiliated to Tel Aviv University, Technion and Israel Space Agency, Israel; (3) Dipartimento di Fisica "E. Amaldi", Università "Roma Tre", Rome, Italy (4) Physics Department, University of Athens, Greece; levpust@ccsg.tau.ac.il

We present developing of methods (e.g., Dorman et al., 1995, 1999) to forecast on the basis of NM hourly on-line data (as well as on-line muon telescopes hourly data from different directions) geomagnetic storms of scales G5 (3-hour index of geomagnetic activity $K_p=9$), G4 ($K_p=8$) and G3 ($K_p=7$) (according to NOAA Space Weather Scales). These geomagnetic storms are dangerous for people technology and health (influence on power systems, on spacecraft operations, on HF radio-communications and others). We show that for especially dangerous geomagnetic storms can be used global-spectrographic method if on-line will be available 35-40 NM and muon telescopes. In this case for each hour can be determined CR anisotropy vector, and the specifically behavior of this vector before SC of geomagnetic storms G5, G4 or G3 (according to NOAA Space Weather Scales) can be used as important factor for forecast. The second factor what can be used for SC forecast is specifically behavior of CR density (CR intensity) for about 30-15 hours before SC (caused mainly by galactic CR particles acceleration during interaction with shock wave moved from the Sun). We demonstrate developing methods on several examples of major geomagnetic storms.

REFERENCES:

Dorman L.I., Villoresi G., Belov A.V., Eroshenko E.A., Iucci N., Yanke V.G., Yudakhin K.F., Bavassano B., Ptitsyna N.G., Tyasto M.I., "Cosmic-ray forecasting features for big Forbush-decreases". Nuclear Physics B, Vol. 49A, pp. 136- (1995).
L.I.Dorman, N.Iucci, N.G.Ptitsyna, G.Villoresi, 1999. "Cosmic ray Forbush-decrease as indicators of space dangerous phenomenon and possible use of cosmic ray data for their prediction", Proc. of 26-th Intern. Cosmic Ray Conference, Salt Lake City, Vol. 6, p.