

RECENT STATUS OF YBJ NEUTRON MONITOR OBSERVATION

H. Miyasaka (1), S. Shimoda (1), Y. Yamada (1), E. Sakamoto (1), K. Munakata (2), S.W.Cui (3), H.H.He (3), H.B.Hu (3), C.L.Lan (3), H.Lu (3), G.X.Luo (3), P.R.Shen (3), Y.H.Tan (3), H.Wang (3), C.Y.Wu (3), H.M.Zhang (3), J.L.Zhang (3), H.W.Guo (4), Labaciren (4), X.R.Meng (4), A.F.Yuan (4), Zhaxisangzhu (4), Zhaxiciren (4), J.Mu (5), X.C.Yang (5), G.M.Le (6) and Z.H.Ye (6)

(1) RIKEN(The Institute of Physical and Chemical Research), Wako, Saitama, Japan, (2) Department of Physics, Shinshu University, Nagano, Japan, (3) The Laboratory of Cosmic Ray and High Energy Physics, IHEP, BeiJing , China, (4) Department of Mathematics and Physics, Tibet University, Lhasa, China, (5) Department of Physics, YunNan University, KunMing, China, (6) Center for Space Science and Applied Research, CAS, Beijing, China.

`miyasaka@riken.go.jp`

The solar neutron observation at Yangbajing (30.11°N 90.53°E, 4300m above sea level, Cutoff rigidity 14.1GV), Tibet has started on October 1998. Now we have almost 3 years data during the solar maximum period. This station performed very high-counts rate (1.07×10^7 counts per hour) with 28 NM-64 counters and high time resolution. We will describe the recent status of intensity variation related to the solar activity and the features of the short period intensity variation.