

SHOCK WAVE ACCELERATION EFFECTS IN THE RADIATION ENVIRONMENT OF THE EARLY SOLAR SYSTEM

L. L. Kashkarov and G. K. Ustinova

Institute of Geochemistry and Analytical Chemistry, RAS, Moscow 117975 Russia.

Observations of active star-forming regions in our Galaxy testify to high intensity radiation in those regions, to its high rigidity, and to the possible enrichment of the spectrum with heavier nuclei. All the effects could be caused by acceleration of particles during the propagation of strong shock waves. The peculiar features of irradiation in the shock wave acceleration conditions led to the change of isotope production rates and to the formation of the specific isotopic anomalies in the matter. Such isotopic anomalies could be conserved in the matter of the solar system, which makes it possible to study the radiation environment at the early solar system stage.

Modelling such processes and analysing the isotopic anomalies in chondrites testify, indeed, to the high intensity and rigidity of irradiation, at least, in some condensation reservoirs of the primordial matter. The average radiation conditions over $\leq 10^7$ years before the termination of accretion are characterized by the average integral proton flux $I_p(> 15MeV) = 1.62 \cdot 10^{19} \text{ cm}^{-2}$ (at $\gamma = 2.5$ and $\alpha/p = 0.0102$). However, during some extreme processes, accompanied by the strong shock waves, the power-law energy spectrum became flatter (up to $\gamma \sim 1.2$), and the integral fluxes $I_p(> E_0)$ were enhanced by 1-2 orders of magnitude. It is just the fact that follows from the data on extinct radionuclides in carbonaceous chondrites. It is possible that the matter of the carbonaceous chondrites was condensed in the explosion shell of the last supernova.

The knowledge of the heavy irradiation component at the stage of the active sun is available due to the tracks of VH-group nuclei ($23 < Z < 28$) observed in silicate minerals of some low-metamorphised ordinary chondrites. Several rare grains ($< 1\%$ of ~ 300) are found in which track gradients from single events of irradiation are revealed. As a general feature, the profiles of these gradients are flatter ($g \sim 0.7 - 2.0$) than those for VH-nuclei of the solar cosmic rays ($g \sim 3.0$), corresponding to the more rigid irradiation. The absence of superposition of the irradiation events characterizes these tracks, i.e. they were produced due to some local single events of the flare pulse type, caused, perhaps, by the reconnection of the magnetic fields in the plasma of the protosolar winds. Obviously, the frequency of the irradiation events was less than the velocity of the matter accretion. The average integral flux of VH-nuclei can be estimated as $I_{VH}(> 15MeV/nucleon) \sim 2.6 \cdot 10^{16} \text{ cm}^{-2}$ at $\gamma \sim 2.5$, its increase being possible by 1-2 orders of magnitude if $\gamma \rightarrow 1$.