

## Neon galactic cosmic ray isotopic abundances: Comparison with Wolf-Rayet star models and meteoritic abundances

W. R. Binns<sup>1</sup>, M. E. Wiedenbeck<sup>2</sup>, E. R. Christian<sup>3</sup>, A. C. Cummings<sup>4</sup>, P. L. Hink<sup>1</sup>, M. H. Israel<sup>1</sup>, R. A. Leske<sup>4</sup>, R. A. Mewaldt<sup>4</sup>, E. C. Stone<sup>4</sup>, T. T. von Rosenvinge<sup>3</sup>, and N. E. Yanasak<sup>4</sup>

<sup>1</sup>Washington University, St. Louis, MO 63130 USA

<sup>2</sup>Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA 91109 USA

<sup>3</sup>NASA/Goddard Space Flight Center, Greenbelt, MD 20771 USA

<sup>4</sup>California Institute of Technology, Pasadena, CA 91125 USA

**Abstract.** Measurements of the neon isotopic abundances by the ACE-CRIS experiment have been obtained in seven energy intervals over the energy range of  $\sim 80 \leq E \leq 280$  MeV/nucleon. The  $^{22}\text{Ne}/^{20}\text{Ne}$  source ratio is derived using the measured  $^{21}\text{Ne}$  abundance as a tracer of secondary production of the neon isotopes. We find that the  $^{22}\text{Ne}/^{20}\text{Ne}$  abundance ratio at the cosmic-ray source is a factor of  $5.0 \pm 0.2$  greater than for the solar wind. The GCR  $^{22}\text{Ne}/^{20}\text{Ne}$  ratio is also shown to be considerably larger than that found in

anomalous cosmic rays, solar energetic particles, and most meteoritic samples of matter. Recent two-component Wolf-Rayet and supernovae models in which GCRs at Earth preferentially sample material from the galactic interior provide predictions for the  $^{22}\text{Ne}/^{20}\text{Ne}$  ratio and other isotope ratios. We will compare the CRIS neon, magnesium, silicon, and iron isotopic source abundance ratios with predictions from these models.