

A search ^7Be in solar energetic particle events

E. R. Christian¹, C. M. S. Cohen², A. C. Cummings², R. A. Leske², R. A. Mewaldt², E. C. Stone², T. T. von Rosenvinge¹, and M. E. Wiedenbeck³

¹NASA/Goddard Space Flight Center

¹California Institute of Technology

¹Jet Propulsion Laboratory

Abstract. The LDEF spacecraft measured an unexpectedly high level of ^7Be , presumably swept up from the residual atmosphere. However, it is unclear whether the atmospheric ^7Be is due to particles originating in nuclear reactions at the Sun, or due to secondary fragments from collisions between energetic particles (solar and galactic) and the atmosphere. The Solar Isotope Spectrometer (SIS) on ACE has been measuring solar energetic particles since launch in 1997. We will present the results of a search for solar ^7Be above about 5 MeV/nucleon with the SIS instrument.

1 Introduction

After its recovery in January 1990, the Long Duration Exposure Facility (LDEF) was discovered to have collected a large amount of the radioactive isotope beryllium-7 (half-life 53.3 days) on its forward-facing surfaces (Fishman *et al.*, 1991; Phillips *et al.*, 1991). The amount of ^7Be measured required a concentration at LDEF altitude (> 320 km) that was thousands of times higher than has been measured in the atmosphere at 20 – 50 km (Lal and Peters 1967).

In the months before LDEF recovery, the Sun was very active, and it was suggested that the solar energetic particle (SEP) events and flares in this time period contributed to the level of ^7Be recorded (Gregory 1996; Share and Murphy 1997). Two processes have been suggested for the ^7Be increase. Either there was increased fragmentation of atmospheric nitrogen and oxygen by the SEPs (which, in addition to cosmic ray interactions, is believed to be the source of the low altitude beryllium), or there was an enhanced level of ^7Be in the SEP population itself. During flares, ^7Be can be produced by α - α fusion in the solar atmosphere

(Kozlovsky and Ramaty 1974; Kuzhevskij and Lur'e 1997). A recent series of experiments (Phillips *et al.* 2001) indicate that the ^7Be concentration is correlated with SEP fluences, but not with solar activity as measured by the x-ray fluence.

The Solar Isotope Spectrometer (SIS) (Stone *et al.* 1998) on the Advanced Composition Explorer (ACE) has been measuring SEPs beyond the Earth's magnetosphere for more than 3 1/2 years. With its large geometry factor (40 cm² sr) and excellent mass resolution, SIS may be able to observe the flux of ^7Be coming from the flare site. Preliminary analysis of SIS data shows no measurable fluence of ^7Be between about 5 and 70 MeV/nucleon. Upper limits on the ^7Be fluence and its relationship to the measured atmospheric ^7Be will be presented at this conference.

Acknowledgements. This research was supported by the National Aeronautics and Space Administration at the Goddard Space Flight Center, the California Institute of Technology (under grant NAG5-6912), and the Jet Propulsion Laboratory.

References

- Fishman, G. J., *et al.* (1991) *Nature* 349:678-680.
- Gregory, J. C. (1996) *Radiat. Meas.* 26:841-850.
- Kozlovsky, B., and Ramaty, R. (1974) *Astrophys. J.* 191: L43-44.
- Kushevskij, B. M., and Lur'e, V. L. (1997) *J. Moscow Phys. Soc.* 7, 7-16.
- Lal, J., and Peters, B. (1967) *Encyclopedia of Physics* 46/2: 551-612.
- Phillips, G. W., *et al.* (1991) *Adv. In Astronautical Sci.* 74:575-583.
- Phillips, G. W., *et al.* (2001) *Geophys. Res. Lett.* 28:939-942.
- Share, G. H., and Murphy, R. J. (1997) *Astrophys. J.* 485:409-418.
- Stone, E. C., *et al.* (1998) *Space Sci. Rev.* 86:357-408.

Correspondence to:

Christian (erc@cosmicra.gsfc.nasa.gov)