

Properties of EHE gamma-ray initiated showers and their search by AGASA

K. Shinozaki⁴, M. Chikawa², M. Fukushima¹, N. Hayashida¹, K. Honda³, N. Inoue⁴, K. Kadota⁵, F. Kakimoto⁶, K. Kamata⁷, S. Kawaguchi⁸, S. Kawakami⁹, Y. Kawasaki¹⁰, N. Kawasumi¹¹, E. Kusano¹², A.M. Mahrous⁴, K. Mase¹, S. Mizobuchi¹³, Y. Morizane², M. Nagano¹⁴, H. Ohoka¹, S. Ozone¹, N. Sakaki¹⁰, M. Sasaki¹, M. Sasano¹⁵, M. Takeda¹, M. Teshima¹, R. Torii¹, I. Tsushima¹¹, Y. Uchihori¹⁶, K. Yasui², T. Yamamoto¹, S. Yoshida¹, and H. Yoshii¹³

¹Institute for Cosmic Ray Research, University of Tokyo, Chiba 277-8582, Japan

²Research Institute for Science and Technology, Kinki University, Osaka 577-8502, Japan

³Faculty of Engineering, Yamanashi University, Kofu 400-8511, Japan

⁴Department of Physics, Saitama University, Saitama 338-8570, Japan

⁵Faculty of Engineering, Musashi Institute of Technology, Tokyo 158-8557, Japan

⁶Department of Physics, Tokyo Institute of Technology, Tokyo 152-8551, Japan

⁷Nishina Memorial Foundation, Komagome, Tokyo 113-0021, Japan

⁸Faculty of Science and Technology, Hirosaki University, Hirosaki 036-8561, Japan

⁹Department of Physics, Osaka City University, Osaka 558-8585, Japan

¹⁰RIKEN (The Institute of Physical and Chemical Research), Saitama 351-0198, Japan

¹¹Faculty of Education, Yamanashi University, Kofu 400-8510, Japan

¹²KEK, High Energy Accelerator Research Organization, Institute of Particle and Nuclear Studies, Tsukuba 305-0801, Japan

¹³Department of Physics, Ehime University, Matsuyama 790-8577, Japan

¹⁴Department of Space Communication Engineering, Fukui University of Technology, Fukui 910-8505, Japan

¹⁵Communications Research Laboratory, Ministry of Posts and Telecommunications, Tokyo 184-8795, Japan

¹⁶National Institute of Radiological Sciences, Chiba 263-8555, Japan

Abstract.

A fraction of gamma-rays in extremely high energy cosmic rays (EHECRs) is a key parameter to discriminate models of the origin of cosmic rays beyond the predicted Greisen-Zatsepin-Ku'zmin(GZK) cut-off. In this paper, we discuss observable properties of EHE gamma-ray initiated showers by Monte Carlo (MC) studies with AIRES code and QGSJET interaction model and a search for EHE gamma-ray initiated showers with muon data taken by the Akeno Giant Air Shower Array (AGASA). The result of the muon component analysis is explained by a fraction of gamma-ray initiated showers with an upper limit of 32% at 90% confidence level (CL) above 10^{19} eV under an assumption of two component primaries of protons and gamma-rays.

consequently predicted as decay products of supermassive particles (*Eg.*, Berezhinski, *et al.* (1998)) or topological defects (*Eg.*, Berezhinski, *et al.* (1997)), or cascade particles initiated by super high energy neutrinos (*Eg.*, Yoshida, Sigl and Lee (1998)).

In order to search for gamma-ray showers by surface-arrays, a fraction of muons in air shower particles may be a possible observable to distinguish from hadronic showers. Due to an inter-detector spacing of ~ 1 km in AGASA, muon densities at 1000m is used as a mass estimator. At energies of interest, MC studies show that showers of gamma-ray primaries produce a relatively small number of muons, while hadronic primaries leave abundant muons. This means that gamma-ray initiated showers have a muon-poor property. In addition, there are also characteristic effects affecting the electromagnetic component in gamma-ray initiated showers.

In this paper, we will discuss observable properties of gamma-ray initiated showers based on MC studies. The result of a search for EHE gamma-rays and a fraction of gamma-rays in cosmic rays above 10^{19} eV will be presented by detailed analysis of AGASA muon data.

1 Introduction

The cosmic rays with energies beyond the predicted GZK cut-off (Greisen (1966); Zatsepin and Ku'zmin (1966)) have been gradually accumulated. AGASA has observed some ten events above 10^{20} eV in the last 11 years (Sakaki, *et al.* (2001)). So far, many models on their origin have been discussed in literatures. Besides traditional 'Bottom-Up' accelerating mechanisms, some 'Top-Down' scenarios have been proposed. The presence of EHE gamma-rays is

Correspondence to: K. Shinozaki

(kenjikry@crsgm1.crinoue.phy.saitama-u.ac.jp)

2 Experiment

2.1 Apparatus

AGASA is located at Lat. $35^{\circ}47'N$, Long. $138^{\circ}30'E$ and 900m asl. 111 stations with a 2.2m² charged particle detec-

tor cover an area of $\sim 100\text{km}^2$. The triggering requirement is a five-fold coincidence among neighboring detectors in a gate width of $25.6\mu\text{sec}$. The triggering bias is free above $10^{18.8}\text{eV}$. Muon detectors with various areas of $2.8\text{--}10\text{m}^2$ each are in operation at 27 stations in the southern region which are triggered by the charged particle detector at each station. Each muon detector consists of 14–20 proportional counters aligned under shielding materials of 1m-concrete or 30cm-lead. The threshold energy is about 0.5GeV for vertically incident showers. The muon density in this work is determined by the so-called ‘on-off’ method. Details are referred to Chiba, *et al.* (1992) and Ohoka, *et al.* (1997).

2.2 Analysis

In this work, events over $10^{18.8}\text{eV}$ are selected by following criteria: 1) Period: December 13, 1995 through December 31, 2000, after unification of AGASA branches; 2) zenith angle $\leq 33.6^\circ$, ($\sec\theta \leq 1.2$); 3) Good fitting of shower geometry; 4) Two or more muon detectors triggered between 800m and 1600m from the core. Criterion 4) is required for the determination of $\rho_\mu(1000)$, the muon density at 1000m from the core, as a mass estimating parameter. $\rho_\mu(1000)$ is obtained by fitting muon densities at that distance range to the empirical lateral distribution function (Hayashida, *et al.* (1995)). The accuracy of $\rho_\mu(1000)$ determination is about 50% on average. The numbers of selected events are 195 and 102 above $10^{18.8}\text{eV}$ and 10^{19}eV , respectively.

In this experiment, we use $S(600)$, the charged particle density at 600m [$1/\text{m}^2$] measured by 5cm-scintillators as an energy estimator (Yoshida, *et al.* (1994)). For hadronic primaries, the uncertainty of the energy estimation is $\sim \pm 30\%$ above 10^{19}eV regardless of the primary species or the interaction model used (Takeda, *et al.* (1998)). The uncertainty for gamma-ray showers will be discussed later.

3 Simulation

We carried out simulation studies using AIRES code (Scuitto (1999)) with QGSJET interaction model (Kalmykov, Ostpcheko and Pavlov (1997)). The zenith angle is fixed to 24.6° , ($\sec\theta = 1.1$). In each combination of primary particles (proton, iron and gamma-ray) and energies ($\log E_0 = 18.0, 18.1, \dots, 21.0$ and $16.0, 16.1, \dots, 21.0$ for gamma-ray primary), 500 showers are simulated and recorded in the library.

EHE gamma-rays above a few times 10^{19}eV may interact with the geomagnetic field to initiate cascades above the atmosphere (McBreen and Lambert (1981)). This effect is examined with the MC code programmed by H.P. Vankov (private communication) which simulates cascades in the geomagnetic field down to the top of the atmosphere. The output is used as an input of AIRES to simulate cascades in the atmosphere.

In order to include the detector fluctuation of measurements, we generated a large number of artificial events us-




Fig. 1. Average $S(600)$ as function of primary energy for $\theta = 24.6^\circ$. The corresponding estimated energy is shown on the right axis. Open circles and squares indicate proton and iron primaries. The solid line shows the gamma-ray. The dotted line shows the conventional energy estimation used in the AGASA Experiment.

ing simulated showers in the library. Shower geometries are sampled from the real data not only to test errors of $\rho_\mu(1000)$ determination and but also to take into account the arrival direction dependence of geomagnetic effects for gamma-ray primaries. The input primary spectrum for each tested primary particle is adjusted to reproduce the observed AGASA cosmic ray spectrum (Hayashida, *et al.* (2000)).

4 Results

4.1 Primary energy estimation

Figure 1 shows average $S(600)$ vs. E_0 relation for proton (open circles), iron (open squares) and gamma-ray (solid curve) primaries for $\theta = 24.6^\circ$. The relation used in AGASA experiment is given by a dotted line and the estimated energy scale is shown on the right axis. The uniform azimuthal distribution is assumed for gamma-ray initiated showers.

For gamma-ray primaries, one needs to note another effect working on shower development; The Landau-Pomeranchuk-Migdal (LPM) effect (Landau and Pomeranchuk (1953); Migdal (1956)). Due to the LPM effect, showers of EHE gamma-ray primaries ‘slowly’ develop and they give smaller $S(600)$ than that of hadronic primaries by 20–30% at energies around 10^{19}eV . This leads to an underestimation of the primary energy of gamma-rays by $S(600)$. This attributes to a suppression of the shower development due to the LPM effect. At much higher energies, gamma-ray primaries mostly interact with the geomagnetic field off the atmosphere and initiate cascading in the magnetic field where no LPM effect or ionization loss works. At the top of the atmosphere, the primary energy is distributed over a bunch of secondary photons and a few electron-positron pairs. The resultant air shower left after this process develops ‘faster’




Fig. 2. $\rho_\mu(1000)$ vs. E_0 relation. Crosses represent individual AGASA events. Muon-less, $\rho_\mu(1000) = 0$ events, are shown in the oval. Dashed, dotted and solid curves indicate upper and lower bounds of 1σ deviation of $\rho_\mu(1000)$ for proton, iron and gamma-ray primaries, respectively.

as a superposition of lower energy subshowers. This transition appears in non-linear $S(600)$ vs. E_0 relation around 5×10^{19} eV because of a competition between LPM and geomagnetic effects on shower development.

4.2 $\rho_\mu(1000)$ vs. E_0 relation

Table 1 summarizes the average $\rho_\mu(1000)$ of showers initiated by proton, iron and gamma-ray with estimated energies of 10^{19} eV, $10^{19.5}$ eV and 10^{20} eV. In the case of gamma-ray primaries, the muon component in the air shower is due to decay products of photonuclear interactions and the average $\rho_\mu(1000)$ is about 20% of that of proton primaries around 10^{19} eV.

In Figure 2, $\rho_\mu(1000)$ vs. E_0 relation is shown. Crosses represent individual events from the experiment. Seven events of $\rho_\mu(1000) = 0$, (called ‘muon-less’ events, hereafter), are indicated at the bottom of the figure. Dashed, dotted and solid curves indicate upper and lower bounds of 1σ deviation of $\rho_\mu(1000)$ for proton, iron and gamma-ray primaries. The general features of the muon component can be explained by hadronic primaries, however, air showers with a small muon content may be a candidate of EHE gamma-ray initiated showers. Above 10^{19} eV, probabilities of ob-

Table 1. $\rho_\mu(1000)[1/m^2]$ at estimated energies of 10^{19} eV, $10^{19.5}$ eV and 10^{20} eV for proton, iron and gamma-ray primaries using the AIREs+QGSJET simulation.

Estimated E_0 [eV]	Proton	Iron	Gamma-ray
$10^{19.0}$	1.5	2.2	0.3
$10^{19.5}$	4.6	6.5	1.0
$10^{20.0}$	11.2	14.0	4.4




Fig. 3. Deduced gamma-ray flux as function of estimated energy. Regions shaded and enclosed by dashed lines indicate upper and lower bounds at a confidence level of 90% for primary combinations of proton+gamma-ray and iron+gamma-ray. Closed circles represent the total cosmic ray flux observed by AGASA (Hayashida, *et al.* (2000)).

serving as muon-less showers are 0.4%, 0.1% and 13% for pure primaries of proton, iron and gamma-ray, respectively. It is noteworthy that four muon-less events above 10^{19} eV were observed against the expected 0.45 events from MC if primaries are mostly protons. The corresponding Poisson chance probability is less than 0.2%. Thus, we can consider these muon-less events as possible candidates for EHE gamma-ray showers.

4.3 Deduction of EHE gamma-ray flux

Figure 3 shows the deduced gamma-ray flux as a function of ‘estimated’ energy. Regions shaded and enclosed by thin dashed lines correspond to the permissible fluxes at 90% CL for assumed two-component primaries of proton+gamma-ray and iron+gamma-ray, respectively. Closed circles indicate the total flux of cosmic rays observed by AGASA (Hayashida, *et al.* (2000)). The flux is a convolution of the AGASA cosmic ray spectrum with the most probable proportion of two components obtained by minimizing χ^2 value for $\rho_\mu(1000)$ distribution in each energy bin of $\Delta \log E_0 = 0.2$. Taking the ambiguities in assumed interaction models into consideration, only an upper limit is given for proton+gamma-ray combination.

As described above, energies of gamma-ray primaries estimated by $S(600)$ is systematically lower than that of hadronic primaries. Therefore the EHE gamma-ray flux reflects that of higher energies which depend on a spectral index of gamma-ray energy primaries. The fraction of EHE gamma-rays above 10^{19} eV yields an upper limit of 32% at 90% CL for the proton+gamma-ray combination. For the iron+gamma-ray, the fraction is $31^{+16}_{-11}\%$. Super- and sub-

scripts denote upper and lower bounds at 90% CL. Corresponding intensities are $< 5.3 \times 10^{-15} [1/m^2 \text{ s sr}]$ and $5.1_{-1.1}^{+2.6} \times 10^{-15} [1/m^2 \text{ s sr}]$, respectively. An upper limit of the EHE gamma-ray intensity at $10^{19.5} \text{eV}$ is estimated to be 50% for that of proton+gamma-ray combination.

5 Discussion

5.1 Individual Super-GZK events

Super-GZK events observed by AGASA are discussed in this Proceedings (Sakaki, *et al.* (2001); Takeda, *et al.* (2001)). The significant event clusters above $4 \times 10^{19} \text{eV}$ events are reported which suggest a possible presence of neutral particles from compact sources. As seen in Figure 2, such highlighted Super-GZK events seem to be explained by hadronic showers. Therefore, there is no explicit indication of gamma-ray primaries at those energies.

5.2 Dependence on interaction models

We have tested observable properties of EHE gamma-ray initiated showers using MC. For an experimental threshold energy ($0.5 \text{GeV} \times \sec \theta$), the observed muon component is derived from interactions at very high energies. Under the photonuclear interaction modeled in EGS4 package (Nelson, Hirayama and Rogers (1985)), which is employed by AIRES, there are less muons in gamma-ray showers component than in hadronic ones. Note that there are claims on a contents of photoproduced muons in gamma-ray initiated showers (*Eg.* Aharonian, Kanevsky and Sahakian (1991)). The muon content in hadronic showers also depends upon interaction models used. Therefore a detailed test of interaction models is required by means of both simulations and experiments in future.

5.3 Fraction of EHE gamma-rays

A fraction of gamma-rays around 10^{19}eV is a realistic threshold energy to test models of the origin of the highest energy cosmic rays with present data. The deduced intensity at an estimated energy depends on that at a higher primary energy. The enhancement factor to convert to primary energy is 20% for the typical gamma-ray differential spectral index of ~ -1.5 predicted by Top-Down scenarios. Under Top-Down scenarios, primaries of heavy nuclei are not expected. The intensity for the proton+gamma-ray primary combination gives a relevant constraint to such models. As mentioned above, the fraction upper bound of EHE gamma-ray is 32% above a primary energy $\sim 10^{19} \text{eV}$. Since Top-Down scenarios predict much lower gamma-ray flux around 10^{19}eV than the present upper limit, this result gives no strong constraints against such models.

6 Conclusion

We have searched for EHE gamma-ray initiated showers with AGASA muon data. Assuming QGSJET interaction models, $\rho_\mu(1000)$ distribution obtained by AGASA is explained by a mixture of hadron and gamma-ray primaries. The contribution of gamma-ray primaries is permitted up to 32% at 90%CL if hadronic primaries are all protons. Hence typical Top-Down models may not be ruled out by the present results.

Acknowledgements. We are grateful to the people in villages and towns, Akeno-mura, Nirasaki-shi, Sutama-cho, Nagasaka-cho, Takane-cho and Ohoizumi-mura for their kind cooperation. The authors are indebted to Mr. M. Kawaguchi and other members of Akeno Group for their great help in the operation and the maintenance of AGASA detectors. K. Shinozaki wishes to acknowledge Prof. H.P. Vankov (Institute for Nuclear Research and Nuclear Energy, Bulgaria) for allowing to use his simulation code and valuable discussions. This work is supported in part by grants in aid #12304012 and #11691117 for the scientific research of JSPS (Japan Society for the Promotion of Science).

References

- Aharonian, F.A., Kanevsky, B.L. and Sahakian, V.A., J.Phys. G: Nucl. Part. Phys. 17 (1991) 1909.
- Berezinski, V., Kachelrieß, M. and Vilenkin, A., Phys. Rev. Lett. 79 (1997) 4302.
- Berezinski, V., Blasi, P., Vilenkin, A., Phys. Rev. D58 (1998) 107.
- Chiba, N., *et al.*, Nucl. Instrum. Meth. A311, (1992) 338.
- Greisen, K., Phys. Rev. Lett. 16, (1966) 748.
- Hayashida, N., *et al.*, J.Phys. G: Nucl. Phys. 21 (1995) 1101.
- Hayashida, N., *et al.*, preprint astro-ph/0008102, (2000).
- Kalmykov, N.N., Ostapcheko, S.S. & Pavlov, A.I., Nucl. Phys. B52 (1997) 17.
- Landau, L.D. and Pomeranchuk, I.J., Dok. Akad. Nauk. SSSR, 92 (1953) 535.;
- McBreen, B. and Lambert, C.J., Phys. Rev. D24 (1981). 2536. See also Kasahara, K., in Proc. of the Intern. Symp. on Extremely High Energy Cosmic Rays, ed. by M. Nagano (Institute for Cosmic Ray Research, University of Tokyo, Tokyo), (1996) 221; Stanev, T. and Vankov, H.P., Phys. Rev.D55 (1997) 1365.
- Migdal, A.B., Phys. Rev. 103 (1956) 1811.
- Nelson, W.R., Hirayama, H. and Rogers, D.W.O., Report SLAC 265 (1985), Stanford Linear Accelerator Center.
- Ohoka, H., *et al.*, Nucl. Instrum. Meth. A385 (1995) 268.
- Sakaki, N., *et al.*, two papers in this Proceeding, (2001).
- Scuitto, S.J., preprint astro-ph/9911331, (1999).
- Takeda M., *et al.*, Phys. Rev. Lett. 81, (1998) 1163.
- Takeda, M., *et al.*, in this Proceedings, (2001).
- Vankov, H.P. (private communication).
- Yoshida, S., *et al.*, J.Phys. G: Nucl. Phys. 20 (1994) 651.
- Yoshida, S., Sigl, G. and Lee, S., Phys. Rev. Lett. 81 (1998) 5505.
- Zatsepin, Z.T. and Ku'zmin, V.A., Pisma Zh. Eksp. Teor. Fiz. 4, (1966) 144.